

Strong Chaos in N-body problem and Microcanonical Thermodynamics of collisionless self gravitating systems.

Piero Cipriani*

Dipartimento di Fisica - Università di Roma "La Sapienza" - P.le A.Moro, 2 - 00185 ROMA (Italia)

Marco Pettini†

Osservatorio Astrofisico di Arcetri L.go E.Fermi, 5 - 50125 FIRENZE (Italia)

February 1, 2008

Abstract. The dynamical justifications which lie at the basis of an *effective* Statistical Mechanics for self gravitating systems are formulated, analyzing some among the well known obstacles thought to prevent a *rigorous* Statistical treatment. It is shown that N-body gravitational systems satisfy a *strong chaos* criterion, so supporting the assumption of an increasingly uniform spreading of orbits over the constant energy surface, *i.e.*, the asymptotic evolution towards a *microcanonical* distribution. We then focus on the necessary conditions for the *equivalence of statistical ensembles* and remark that this equivalence is broken for any N-body system whose interaction has a range comparable with its spatial extension. Once realized that the obstacles originate from the long range nature of the Newtonian interaction, leading to the non-extensivity of canonical and grand-canonical thermodynamic potentials, we show that instead a suitably generalized *microcanonical ensemble* constitutes an *orthode*, *i.e.*, a reliable framework for a *correct* Thermodynamics. Within this setting we use the thermodynamic relations and find consistent definitions for Entropy (which turns out to be *extensive*, despite the non stable nature of the interaction), Temperature and Heat Capacity. Then, a *Second Law-like* criterion is used to select the hierarchy of *secular equilibria* describing, for any finite time, the macroscopic behaviour of self gravitating systems.

Keywords: Gravitational Physics, N-body Systems, Statistical Mechanics

1. Introduction

Although is the most *ancient* among the known *fundamental* interactions, Gravity seems nevertheless to be the most reluctant to be included coherently within the theoretic frameworks describing the behaviour of *microscopic* and *macroscopic* many body systems, respectively.

Indeed, in addition to the well known difficulties encountered by the attempts for a coherent formulation of a Quantum Theory of Gravity,

* Permanent address: **C.S.S.** - via E.Nardi, 14/16 - 02047 - Poggio Mirteto (RI) Italia - Electronic address: piero.cipriani@roma1.infn.it or cipriani@icra.it

† Also I.N.F.N., Sezione di Firenze - Electronic address: pettini@arcetri.astro.it



there are the less *popular*, though equally relevant from a conceptual viewpoint, obstacles preventing a *rigorous* Statistical Mechanics of collisionless self gravitating N-body systems (1).

This is due to the fact that the Hamiltonian function describing the gravitational interaction between N point particles of mass m_i

$$H(q, p) = \sum_{i=1}^N \frac{\mathbf{p}_i^2}{2m_i} - \frac{1}{2} \sum_{i=1}^N \sum_{j \neq i}^N \frac{Gm_i m_j}{r_{ij}}, \quad \text{where } r_{ij} \equiv \|\mathbf{q}_j - \mathbf{q}_i\| \quad (1)$$

does not satisfy the *stability* property, which is a prerequisite for a self consistent Statistical Mechanical description and for the proofs of most of the theorems assuring the equivalence of the Statistical ensembles in the so-called *Thermodynamic limit*. The lack of ensemble equivalence for N-body self gravitating systems (SGS), leads naturally to the issue of determining which, if any, among the many ensembles complies with the *orthodic* condition (2).

Notwithstanding the hindrances for a Statistical Mechanics of gravitational systems are known since many decades, many efforts have been devoted to overcome them, in the attempt to formulate, at least, an *effective* Statistical description of collisionless N-body SGS (3; 4; 5; 6).

The astrophysical motivations for doing so are well evident looking at the otherwise inexplicable morphological, structural and dynamical similarities among *hot* stellar systems observed in the sky. The *qualitative* homogeneities of the luminosity curves and of velocity dispersion profiles, though with obvious *quantitative* differences, suggest unavoidably the existence of one or more relaxation processes leading globular clusters and elliptical galaxies towards an equilibrium state which is qualitatively independent on the global parameters of the systems and on the class of initial conditions of these astronomical objects at the formation era.

Due to the peculiar nature of Gravity, the relaxation mechanisms (e.g., binary *collisions*) at work in usual *laboratory* many body systems turn out to be unable to justify such a state of (*quasi*) equilibrium, because the corresponding *relaxation time* predicted is comparable to the Hubble time for globular clusters, and several orders of magnitude larger for elliptical galaxies.

More than thirty years ago, Lynden-Bell, Saslaw and others (7; 8) first realized that the infinite range of Newtonian interaction could support relaxation mechanisms related to *collective effects*, in turn due to global, almost coherent, oscillations of the gravitational field, able to change appreciably the energies of the particles (*i.e.*, the stars). However, the general relevance of such a mechanism has been revised and even criticized to some extent, (9; 10) and it is by now agreed that,

though useful to understand some aspects of the initial global evolution of gravitational systems, it alone cannot be sufficient to explain the *universal* light and velocity profiles observed.

The lack of a satisfactory Statistical Mechanics for SGS made problematic obviously any attempt to formulate a Thermodynamics for them. Among the various *peculiarities* of gravitational interaction (discussed in the sections below), those implying the *non extensivity* of (dynamic and) thermodynamic potentials constitute the main obstacles to the definition of thermodynamic quantities and, even more, to derive *standard* relationships among them.

A major piece of evidence of the very peculiar Thermodynamics proper to SGS is known since many decades to astronomers, but it went to know a wide *popularity* in the Physics community only when Thirring *rediscovered* this apparent *paradox* (11), which was already investigated rather deeply some years before by Antonov (12) and by Lynden-Bell and Wood (13). The *problematic issue* emerges when we recall from one side the well known (to astronomers) property of (bound) SGS of having *negative heat capacities*, and from the other the expression of the heat capacity as it can be easily derived within the Canonical ensemble (see, e.g., (14), §8.2):

$$\frac{\partial U}{\partial \beta} + \langle (H - U)^2 \rangle = 0 , \quad (2)$$

where $U \equiv \langle H \rangle$ is the average value of the energy of the system in the canonical ensemble, $\beta = 1/k_B T$ is the (inverse) temperature parameter of the canonical distribution and k_B is the Boltzmann constant. Written in terms of the heat capacity C_V , equation (2) reads

$$C_V = \frac{1}{k_B T^2} (\langle H^2 \rangle - \langle H \rangle^2) \geq 0 . \quad (3)$$

In the recent years, however, it has been realized that such a paradox is not a peculiarity exclusive to SGS, but it is instead a common feature of *small systems*, *i.e.*, systems whose extension is smaller than (or comparable to) the range of the governing interaction (15; 16; 17). For these systems it is immediately realized that *extensivity* property breaks down. However these systems possess a well defined, though in some respects peculiar, thermodynamical behaviour, and so it turns out as well that such a property is not necessarily relevant to allow for a description of the macroscopic, *quasi-thermodynamic* behaviour of these systems, as discussed in more detail below, where we will analyze also why this feature obstructed to some extent the formulation of a Thermodynamic framework for SGS.

An immediate consequence of the lack of extensivity is the breakdown of the *ensemble equivalence*, which is another milestone of the *rigorous* statistical mechanical foundation of the thermodynamic formalism. However, the phenomenology of non extensive systems is currently investigated from theoretical, numerical and also experimental viewpoints, and many studies nowadays focus on the dynamical, statistical and thermodynamic properties of Hamiltonian systems characterized by ensemble inequivalence (15; 17; 18).

According to the considerations above, the attempt to formulate a *coherent* thermodynamic description of SGS appear to be feasible, starting from the basic assumptions which lie at the grounds of the Statistical description of many degrees of freedom dynamical systems, though paying due attention to every possible source of internal inconsistency.

Then, going back to Dynamics and Statistical Mechanics, we first investigate the requisites to be fulfilled by the Dynamics, in order to justify a statistical description of macroscopic properties of N-body systems. This goal is accomplished using some concepts of Ergodic theory of (many degrees of freedom) dynamical systems, and in particular we adopt the *Geometric Description* of Dynamics, which allows to relate the chaotic properties of motions to the geometrical and topological properties of the dynamical manifolds associated to the system, see, e.g., (19; 20; 21; 22; 23; 24).

Once found convincing indications that a self consistent *effective* Statistical Mechanics of SGS can be formulated¹, we then face the issue of the ensemble inequivalence and argue that the Classical *Microcanonical ensemble* alone is suitable for a reliable and coherent description of the Thermodynamics of non extensive systems like gravitational N-body ones. The criteria according to which the Microcanonical distribution over the hypersurface $\Sigma_{\mathcal{E}}$ of the full $6N$ -dimensional phase space Γ , acquires such a privileged role over other classical ensembles are analyzed and discussed, and it is shown how such a choice allows to derive a number of *physically reasonable* results on the macroscopic properties of N-body gravitating systems as well as unambiguous (re-)definitions of thermodynamic quantities and relations.

As a by-product of the above analysis, we also derive and discuss some among the many non trivial relationships existing between the statistical properties² of dynamics over the constant energy hyper-

¹ Though with some peculiarities which are the statistical counterparts of the dynamical unique features of gravitational interaction.

² As, for example, ergodicity of dynamics in Γ space or even more strong features, like *mixing*.

surface $\Sigma_{\mathcal{E}}$, the consequent statistical mechanical description, and the customary *reduced* description in terms of a *distribution function* (DF) over the single particle six dimensional μ -phase space, depending only, according to Jeans theorem (6), on the isolating integrals of motion admitted by the *smoothed* potential in the three dimensional configuration space. This theorem indeed is at the grounds of most of the works devoted to the construction of self-consistent models of stellar systems over the last four decades, and only recently it has been realized that its relevance for stellar dynamical modelling, at least in the customary *strong formulation* sketched above, has been somewhat overemphasized (25). We outline here the conditions to be satisfied by the full dynamics in Γ_{6N} -space in order to allow for the *reduction process* leading to a hydrodynamic (or kinetic) description in terms of a distribution function over the μ -space. This represents a first step along the path of a critical reconsideration of the use of the single particle DF as a tool to derive the equilibrium properties of stellar systems through *maximum entropy* like arguments based on it (26). A deeper discussion of this topic is however beyond the scope of this paper and will be the subject of forthcoming works.

The paper is organised as follows: in section 2 we study the chaotic properties of many degrees of freedom systems, with particular emphasis on the features of the dynamics of SGS and recall the usuals indicators adopted to quantify the degree of stochasticity of orbits in phase space. In order to relate Dynamics to the structure of the ambient space, we adopt the terminology and the tools of the Geometric Description of Dynamics, which provides an elegant and effective framework to characterize the intrinsic properties of dynamical systems.

To support our argumentations we recall some of the results obtained within this framework and reinterpret them in the perspective of the present work, devoted to propose thermodynamical description of SGS, justified on the grounds of the statistical features of dynamics and of the geometric properties of the dynamical manifolds. In section 3 we briefly introduce some concepts from the Ergodic theory of Dynamical Systems and use them to classify the statistical properties of the flow associated to the N-body dynamics, arguing for the possibility of a statistical description of their macroscopic properties on any finite scale of times. In section 4 we review the issues related to the non equivalence of ensembles, then we discuss why the classical Microcanonical ensemble is the only legitimate to describe coherently the "*Thermodynamics*" of SGS; we then present some consequences of the thermodynamic relations derived, analyzing also the various inconsistencies encountered attempting to describe the system within

the *canonical* ensemble. Section 5 presents a brief discussion of the results, the *peculiarities* of the Thermodynamics of systems without *Thermodynamic limit*, a heuristic discussion on the concept of *finite time equilibrium* and concludes outlining the work in progress along the line here reported, possible extensions of the formalism and the improvements to the results presented which can be obtained with further studies and more extensive numerical work.

2. Chaotic behaviour and Ergodic properties of Dynamics of gravitational N-body systems.

Since the first *rediscoveries*³ of the ubiquity of Chaos in nonlinear dynamical systems in the 60's, (28), it was realized that unpredictability implied by the exponential sensitivity to small changes in initial conditions could be the ingredient needed to justify a more or less *physical* reformulation of the *ergodic hypothesis*, able to explain the approach to a stationary equilibrium state of most non integrable many degrees of freedom (MDoF) Dynamical Systems. To make a long story short, it is by now generally agreed that the presence of Chaos entails a system to forget the detailed memory of initial conditions, though some *imprinting* related to few global constraints or *quasi conserved* quantities could survive during the dynamical evolution, leading to a slow decay in time of specific correlation functions. The sufficient conditions implying irreversible behaviour and then able to justify a statistical mechanical description of macroscopic properties have been mathematically demonstrated only for very special (and rather abstract) classes of models, however there is a widespread agreement that, from a physical viewpoint, most MDoF nonlinear dynamical systems are indeed *chaotic enough* to relax at least to a stationary (meta-) equilibrium state.

2.1. CHAOS, ERGODICITY AND MIXING.

For few dimensional systems Chaos is not sufficient to guarantee the efficiency of the approach to equilibrium, because diffusion in phase space can be inhibited by the fractal structure of deformed (or partially destroyed) invariant surfaces describing the *integrable limit*, *i.e.*, by the *Cantori* originated by the disappearance of KAM-tori (29). Even above the KAM threshold, the exponentially long times associated to

³ Most of the conceptual implications of dynamical instability as well as the realization that such a phenomenology should be the rule rather than the exception in the behavior of nonlinear (*i.e.*, generic) dynamical systems were indeed already present in the Poincaré work (27) more than a century ago!

the Nekhoroshev regime⁴(30), the relaxation processes are practically *frozen* and from a physical viewpoint the approach to equilibrium will not take place at any finite time (31).

However, both KAM and Nekhoroshev thresholds depends strongly on the dimensionality of phase space⁵, decreasing roughly exponentially with the number of degrees of freedom. Together with the Poincaré-Fermi (32) theorem on the non existence of analytic integrals for non-linear systems with three or more degrees of freedom, the vanishing of the integrability or quasi integrability thresholds, suggests that MDoF chaotic systems should attain the *asymptotic equilibrium states*.

However, from a physical viewpoint, it is also important that the approach to such states be fast enough, that is, it is required that the time averages of relevant dynamical macroscopic quantities converge to corresponding phase averages in finite times and not only asymptotically. Rephrased in terms of Statistical theory of Dynamical Systems, this amounts to say that dynamics leads any finite volume in phase space to spread out uniformly over the entire region accessible to motion. Formally, in addition to the *ergodic* property:

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T F[\mathbf{q}(t), \mathbf{p}(t)] dt = \frac{1}{\mathcal{M}(\Gamma)} \int_{\Gamma} F(\mathbf{q}, \mathbf{p}) \mu(d\mathbf{x}) , \quad (4)$$

the dynamical evolution must also satisfy the stronger condition known as *mixing*:

$$\lim_{T \rightarrow \infty} \int_{\Gamma} F[\mathbf{q}(t), \mathbf{p}(t)] G[\mathbf{q}, \mathbf{p}] \mu(d\mathbf{x}) = \left(\int_{\Gamma} F(\mathbf{q}, \mathbf{p}) \mu(d\mathbf{x}) \right) \cdot \left(\int_{\Gamma} G(\mathbf{q}, \mathbf{p}) \mu(d\mathbf{x}) \right) , \quad (5)$$

where F and G are arbitrary functions of dynamical variables and $\mu(d\mathbf{x})$ is the invariant measure over Γ ⁶.

As above, the *mixing* property has been rigorously proved only for restricted classes of abstract dynamical systems, *e.g.*, (29), though many theoretical and numerical works (33) now support strongly the belief that even more realistic models of many body physical systems

⁴ During which the actions are almost exactly conserved and dynamical trajectories are trapped between confining surfaces.

⁵ Throughout the paper we indicate with $\mathcal{N} = N \cdot d$, the number of degrees of freedom of a dynamical system, N being the number of particles and d the dimensionality of the single particle configuration space. For Hamiltonian systems then the phase space Γ has dimension $2\mathcal{N}$ and the constant energy hypersurface $\Sigma_{\mathcal{E}}$ has clearly dimension $2\mathcal{N} - 1$.

⁶ What is relevant from the point of view of Physics is that equation 5 be satisfied at least by macroscopic *relevant* functions F and not necessarily for *any* pair F and G , as instead is required in the mathematical definition of *mixing*. In the case of Hamiltonian systems the invariant measure is the Liouville measure.

possess *good* statistical properties and approach the equilibrium state on physically reasonable time scales; moreover, these statistical relaxation times bear close correlations with the orbit diffusion rates in phase space, and also with Lyapunov exponents describing the degree of chaoticity.

The above evidences have been given a formal expression recently through the formulation of the so-called *Chaotic Hypothesis* (34), which puts on the same level, for what concerns the statistical description of macroscopic properties, chaotic many degrees of freedom dynamical systems and Anosov systems, which are, among the *abstract* models, those characterized by the strongest statistical properties (together with *Axiom A* and *K-systems*, see, e.g., (35))⁷.

Before to proceed further, it is necessary to remark that all the above discussion about the KAM and Nekhoroshev thresholds is of fundamental relevance to single out the sources of Chaos in MDoF dynamical systems and for the understanding of the relationships between the instability of the trajectories and the modifications occurring in the structure of phase space. Nevertheless, most of the tools and concepts developed within that framework are of little use for the characterization of the dynamical properties of SGS in an arbitrary state, because there is nothing indicating the existence of any integrable limit for gravitational N-body systems; *i.e.*, for them, there is no hope to undertake a sensible perturbative approach to study the onset of Chaos, at least as long as $\ln N \geq 1$. This amounts to say that the investigation of dynamical instability in SGS must resort to methods and tools pertaining more to the theory of strongly chaotic dynamical systems rather than to Perturbation Theory, whose approaches are suited for nearly integrable systems.

2.2. GEOMETRY OF DYNAMICS AND CHAOS.

Within the class of systems for which the *mixing* property has been rigorously proved there are the *geodesic flows on manifolds of constant negative curvature* (36), which were studied in the 30's by Hedlund and Hopf. The first to realize the relevance of such a line of research to Statistical Physics was Krylov (37), which studied the consequences of the instability of the geodesic flow associated to the dynamics of an hard sphere gas, predicting that the instability leads the system to macroscopically approach the equilibrium state, with a relaxation time which agrees rather well with more recent estimates.

⁷ These models, though maximally chaotic, are, for what concerns the evolution of their statistical properties, almost as simple as integrable systems ((20), chapters 4÷6).

The formal apparatus for the geometric transcription of the dynamics of natural systems was developed independently by many authors and found a beautiful presentation in the Synge's article "*On the Geometry of Dynamics*" (38). Although completely unaware of the relevance of exponential instability, he there discussed extensively the stability of geodesic flows of dynamical interest.

The fundamental argument able to relate geodesic instability and the entailed chaotic dynamics from one side, to the fulfillment of the *mixing* property on the other, goes through the proof that the dynamical manifold is characterised by almost everywhere negative sectional curvatures. If this is the case, then the *Instability time*, as measured by a suitable generalization of the (inverse) Lyapunov exponent, is closely related to the Relaxation time, giving the scale over which correlations decay and the memory of the initial state is lost. Then the point has been moved from a direct proof of mixing to the study of the sign of the sectional curvatures of the manifold.

We refer for the details of the geometrical transcription of dynamics of Hamiltonian and Lagrangian systems to (24; 22) and references therein, and simply recall that the tool for investigating the stability of the geodesic flow are the *Jacobi-Levi-Civita* equations for geodesic deviation, which, in local coordinates over a suitable manifold read:

$$\frac{\nabla}{ds} \left(\frac{\nabla \delta q^a}{ds} \right) + \mathcal{H}^a{}_c \delta q^c = 0, \quad (a = 1, \dots, \mathcal{N}) \quad (6)$$

where $\delta \mathbf{q}$ represents the perturbation to the flow, whose (possibly) exponential growth is the main signature of instability and depends on the *stability tensor* $\mathcal{H}^a{}_c$, defined *along* a given geodesic (with unitary tangent vector $\mathbf{u}$), through the Riemann curvature tensor, $R^a{}_{bcd}$ of the manifold (with metric g_{ab}), by

$$\mathcal{H}^a{}_c \doteq R^a{}_{bcd} u^b u^d; \quad (7)$$

and ∇/ds is the total (covariant) derivative along the geodesic. From equation (6), an effective ordinary differential equation for the norm of the perturbation can be derived. Defining an unit vector $\boldsymbol{\nu}$ on the unitary tangent bundle of the manifold $\mathbf{M}$, $\mathbf{T}_\mathbf{q}\mathbf{M}$, such that $\delta \mathbf{q} = z \boldsymbol{\nu}$, it is straightforward, (22), to show that the magnitude of the perturbation, z , evolves according to:

$$\frac{d^2 z}{ds^2} = \left(-\mathcal{H}_{ac} \nu^a \nu^c + \left\| \frac{\nabla \boldsymbol{\nu}}{ds} \right\|^2 \right) z \quad (8)$$

which is still an *exact* equation containing all the required informations about the overall instability of the flow. The simplification obtained

going from equation(6) to equation(8) is rather illusory. Indeed, even if we reduced a system of $\mathcal{N}$ differential equations containing covariant derivatives apparently to just one ordinary differential equation, the computational task has been made only a little easier. The problem is that equation (8) still contains the full stability tensor, whose explicit computation requires the knowledge of the $\mathcal{O}(\mathcal{N}^4)$ components of the Riemann tensor. In some particular cases it happens that the manifold is such that the simplification obtained is actually decisive (39), but as expected, this is not true for most geometric transcriptions of realistic models of physical systems.

For MDoF systems, however, in the hypothesis, to be verified *a posteriori*, that the dynamics is *far enough from complete integrability*⁸, it is possible to obtain a *closure* of equation(8), which then reduces to the approximate form:

$$\frac{d^2 z}{ds^2} + \mathcal{K}_{R_u}[\mathbf{q}(s)] z \cong 0 ; \quad (9)$$

where

$$\mathcal{K}_{R_u}[\mathbf{q}(s)] \doteq \frac{\mathcal{H}^a_a}{\mathcal{N}-1} \equiv \frac{\text{Tr } \mathcal{H}}{\mathcal{N}-1} ; \quad (10)$$

is nothing else than the *Ricci curvature* in the direction of the geodesic, equal also to

$$\mathcal{K}_{R_u}[\mathbf{q}(s)] = \frac{R_{ab}u^a u^b}{\mathcal{N}-1} \doteq \frac{\text{Ric}(\mathbf{u})}{\mathcal{N}-1} , \quad (11)$$

where $R_{ac} \doteq R^b_{abc} \equiv g^{bd}R_{abcd}$ is the Ricci tensor.

Referring to the above cited bibliography for a detailed discussion, we simply observe here how, from the effective equation (9) it is easy to understand the classical results on the *chaotic* properties of geodesic flows on manifolds of negative curvature: if the Ricci curvature per degree of freedom $\mathcal{K}_{R_u}[\mathbf{q}(s)]$ is everywhere negative along a geodesic, then the geodesic flow is exponentially unstable. So, a curvature everywhere negative (that is, always, along a geodesic) entails an exponential instability of the flow.

⁸ What is required actually is much less: it is sufficient that the evolution of the many degrees of freedom is not coherent, that is, that there is a spectrum of frequencies wide and far enough from resonances, so that collective periodic oscillations are avoided. In other terms, a good *phase mixing* is sufficient to prevent the persistence of phase correlations among the motions of individual particles; and we recall that phase mixing occurs in integrable systems with MDoF as well! From the geometrical viewpoint, this is enough to imply that the $\mathcal{N}-1$ non vanishing sectional curvatures are locally almost uncorrelated each other.

But this is not the only mechanism leading to Chaos. It is indeed possible to have an exponential growth of the solution of equation (9) also when the sign of the Ricci curvature along a geodesic is fluctuating and even for everywhere positive curvature.

2.2.1. Sources of instability in positive curvature manifolds.

Two further sources of instability are indeed possible for equations of the form (9):

- the onset of exponentially unstable solutions can originate (when $\mathcal{K}_R > 0$) from the mechanism of *parametric resonance*. This phenomenon, well known in the linear case (see, e.g., (40)), occurs for example in the generalization of Mathieu-like equations,

$$\ddot{x} + \Omega^2(t)x = 0, \quad \Omega^2(t) = \Omega_o^2[1 + f(t)], \quad \langle f \rangle_t = 0, \quad (12)$$

when the frequency $\Omega^2(t)$ undergoes a periodic or multiperiodic modulation whose frequencies satisfy suitable resonance conditions with the unperturbed average frequency, Ω_o . If the behaviour of parametric resonance in linear systems is rather elementary, when this phenomenon couples with even small nonlinearities present in the system, the outcome can be very rich and complex (41). The interest in this mechanism resides in the fact that it appears very often when a phenomenological modellization of complex systems is attempted through a reduction process leading to the formulation of a system of *effective equations*, alike equation (9) above. It has been shown (e.g., (22; 23; 24)) that the occurrence of positive curvatures is the rule, rather than the exception, for most geometrical transcription of MDoF dynamical systems. As it is known, nearly all nonlinear systems exhibit nevertheless chaotic behaviours for almost all initial conditions. As every system is characterized by one or more intrinsic periodicities, often related to some kind of collective behaviour⁹, it appears reasonable to investigate the correspondences between the parameter space and the astonishingly rich behaviour disclosed by evolutionary laws apparently so simple. In order to give a taste of the various phenomenologies displayed from an equation with only 1.5 degrees of freedom, we report in figure 1, the behaviour of different solutions

⁹ As examples ranging from Solid State Physics to Astrophysics, we mention the resonances occurring with the lower modes frequencies in a slightly anharmonic crystal, which modulate the dynamics on the largest possible spatial scales, and the overall quasi periodicity of the gravitational field inside a stellar systems undergoing the Violent Relaxation phase, which modulate the background field of existing condensed substructures.

Figure 1. Parametric Resonance in a one degree of freedom nonlinear Lagrangian system with time-dependent potential. Each panel shows the plot of the solution of a single equation with a slightly different choice for the values of the only two free parameters. The behaviour of solutions of the first two panels recall those of the underlying linear model, with the typical damped and resonant evolution, respectively. Other choices of the parameters values originate either quasi periodic solutions, intermittent or steadily *chaotic* patterns.

of a single equation, obtained adding small nonlinearities to (12) and whose two parameters have been varied slightly to obtain the different patterns shown in the panels.

- The second mechanism able to lead to exponential instability of the flow is known as *Stochastic Resonance* and arises instead when the modulation of the frequency is due to stochastic fluctuations around its average value (42), and is typical of either non completely isolated systems, where fluctuations arise from the environmental influence, or, as for parametric resonance, in the case of *effective equations*, describing the evolution of global or collective variables. Now the modulation is not described through deterministic modellization, but instead originates from the fact that the exact values of the parameters (and in particular of the average frequency) are determined by the combined effects of all the *microscopic* constituents of the system and are then influenced by the noise due to finite size effects.

As a general rule (though of a rather heuristic nature), parametric resonance is responsible of the unstable behaviour of few dimensional systems as well as of the dynamical instability occurring in MDoF not very far from integrability and with an overall intrinsic underlying periodicity, whereas, as stated before, stochastic resonance is the typical mechanism promoting the development of Chaos in strongly non integrable MDoF systems; its description becomes more and more accurate as the number of degrees of freedom increases.

It is obviously possible that the two mechanisms operate together, even if in this case it turns out very difficult to separate the effects and even to understand the conditions under which they enhance or damp each other. What is more plausible is that there are regimes in a given model in which either of the two mechanisms dominate over the other. If this is the case, heuristic argumentations suggest that in the intermediate range the two effects substantially do not interact, otherwise a discontinuous transition should be observed¹⁰.

¹⁰ As the two mechanisms cause the onset of instability exciting in a very different way the frequency space (strongly localized the first, with a rather narrow

From the above discussion we expect that the mechanism leading SGS to Chaos to be more closely related to Stochastic rather than parametric resonance, and we will see below that this is indeed the case.

2.2.2. *Geometric properties of the manifold and analytic computation of the maximal Lyapunov exponent.*

In recent years the geometrical description of dynamics has been applied to the investigations of the dynamical behaviour of a large class of Hamiltonian and Lagrangian systems either with few or many degrees of freedom, (see, e.g., (19; 20; 22; 23) and for a comprehensive review and an updated bibliography, (24)).

The *GeometroDynamical Approach* (GDA) applied to Hamiltonian MDoF systems allowed to improve our understanding on many aspects of the characterization, quantification and sources of chaotic behaviour. Among the most remarkable, from the point of view of a concise synthetic characterization of the dynamic instability, it has been possible to use the GDA, in conjunction with the theory of Stochastic Differential Equations (42), to derive an effective analytic formula for the maximal Lyapunov exponent (21), which is obtained simply evaluating average values, fluctuations and autocorrelation time of the *frequency* of the effective equation (9).

If we indicate with $k_R \doteq \langle \mathcal{K}_{Ru}[\mathbf{q}(s)] \rangle$ the average (positive) value of the Ricci curvature per degree of freedom, with $\sigma_{k_R}^2 \doteq \mathcal{N} \sigma_{k_R}^2$ the rescaled variance of its fluctuations¹¹ and with τ_{k_R} the autocorrelation time of the stochastic process describing the time evolution of $\mathcal{K}_{Ru}[\mathbf{q}(s)]$ along any given geodesic, it is possible to analytically compute the average exponential growth rate of the perturbation according to the following formula

$$\lambda_1 = \sqrt{\frac{k_R}{3}} \left(g^{1/3} - g^{-1/3} \right) , \quad (13)$$

where

$$g \doteq \frac{1 + \sqrt{1 + \eta^2}}{\eta} \quad (14)$$

and

$$\eta \doteq \frac{8\sqrt{3}}{9} \frac{k_R^{3/2}}{\sigma_{k_R}^2 \tau_{k_R}} . \quad (15)$$

distribution the second), we expect the effect of the overlap to be in generic cases negligible.

¹¹ For a discussion of the $\mathcal{N}$ rescaling of the variance, see (21).

The importance of equation (13) relies also on the fact that for most many body systems the quantities entering it can be computed using either the standard time averages along numerical trajectories or the *statistical* phase averages over the canonical or microcanonical distribution. The agreement between the results obtained using either of the above procedures is rather satisfactory (20; 21), and numerical simulations along trajectories allow also to check the reliability of the assumptions at the grounds of the use of the theory of stochastic differential equations, first of all the gaussian distribution of the values of the curvature.

In addition to this, the GDA helps to improve the conceptual understanding of the onset of instability in MDoF Hamiltonian systems, showing that, even for the most chaotic N-body systems, the sources of instability rely mostly on the stochastic fluctuations of the curvature rather than on the occurrence of negative values. In particular, for what concerns SGS, it has become clear that dynamical instability of trajectories, originates mainly from *binary encounters*, and that the latter entail *positive fluctuations of the curvature*: the strong chaoticity of N-body systems is mainly due just to the huge amplitude of curvature's fluctuations around the average value, to be compared with what happens in other MDoF Hamiltonian systems, where fluctuations become of the order of the average values only in the strongly chaotic limit.

Moreover, the GDA singles out clearly the different regimes accompanying the approach to equilibrium, as the transition to a *less unstable* regime when virial equilibrium has been attained and the violent relaxation process accomplished; and many other phenomena operating on longer timescales and finer spatial extensions. The details of these issue have been partially discussed in (22) and will be analysed in a deeper detail elsewhere.

2.3. STRONG STOCHASTICITY THRESHOLD AND CHAOTIC HYPOTHESIS.

To the main issue of this paper what is relevant of the above discussion is the evidence of a very strong chaoticity in gravitational N-body systems. Without going into formal details, we present here the results of the application of the above formalism to SGS, and compare them with analogous outcomes related to a chain of anharmonic oscillators¹², in order to characterize the degree of chaoticity present in N-body dynamics. As anticipated above, in MDoF nonlinear systems there are in general various dynamical regimes. Starting from quasi integrable

¹² The well known Fermi-Pasta-Ulam quartic, or FPU- β , model (32).

behaviour for small nonlinearity (or at enough low energy), a complete hierarchy of chaotic regimes is encountered as the departure from integrability increases. The KAM and Nekhoroshev thresholds are very near the *zero* of the perturbation parameter, ε , as soon as the number of degrees of freedom is greater than a few tens, but nevertheless, in several different Hamiltonian systems it is possible to single out a further physically meaningful threshold, separating two different regimes of stochasticity. This dynamical transition has its statistical counterpart in a rather abrupt change in the behaviour of the relaxation time of macroscopic quantities.

Briefly, below such a threshold, let it be ε_c , the rate of increase of the maximal Lyapunov exponent, $\lambda_1(\varepsilon)$, is described by a relatively steep (model dependent) power law and the relaxation times are very long, depending sensitively on the perturbation parameter. Above this threshold, the growth rate of $\lambda_1(\varepsilon)$ slows down and the relaxation times become much shorter and nearly independent of the value of ε (as long as $\varepsilon > \varepsilon_c$) (33).

A very interesting and illuminating feature of the scaling law $\lambda_1(\varepsilon)$ above ε_c ¹³ is such that the dimensionless quantity $\gamma_1 \doteq \lambda_1 \cdot t_D$, introduced in (22), turns out to be virtually independent on the perturbation parameter. This because, above the SST, the maximal Lyapunov exponent scales exactly as the reciprocal of the dynamical time scale. This evidence implies that the diffusion of orbits in phase space proceeds at the fastest rate allowed by the dynamics. This fact has been noticed and briefly discussed in (22), suggests a very elegant scenario for the mechanisms governing the onset and the development of Chaos, and gives a convincing picture of the modifications in the phase space structure accompanying the departure from integrability. The investigation of these issues and of their consequences is however beyond the scope of these pages, and will be discussed elsewhere.

To proceed towards our goal, *i.e.*, a Thermodynamics for SGS, in the figures 2 and 3, we compare the energy dependence¹⁴ of the maximal Lyapunov exponent, computed either using the standard algorithm (43) or the analytic geometrodynamical formula, equation (13).

Figure 2 displays the transition in the scaling behaviour of $\gamma_1(\beta\varepsilon)$ outlined above.

In figure 3 are reported the values of the maximal Lyapunov exponent as a function of the specific energy of the system, for two series of simulations with $N = 400$ and $N = 4000$ bodies. It is evident the power-

¹³ The so called *strong stochasticity threshold*, SST, (19; 33).

¹⁴ To be precise, for the FPU model, the perturbation parameter is the product $\beta\varepsilon$, where β is the anharmonicity constant.

Figure 2. Scaling of the dimensionless chaoticity indicator $\gamma_1 = \lambda_1 t_D$ with the perturbation parameter $\beta\varepsilon$ for a FPU chain, as obtained according to the analytic formula, eq.(13). Open circles refer to the values computed from mean values and fluctuations obtained from phase space averages (21), whereas, starlike symbols with error bars show the values obtained from time averages computed along numerically integrated trajectories (20).

Figure 3. Scaling with the binding energy per particle, ε , of the standard maximal Lyapunov exponent λ_1 and of the dimensionless chaoticity indicator $\gamma_1 = \lambda_1 t_D$ for two sets of gravitational N-body systems, with $N = 400$ (circles) and $N = 4000$ (stars) bodies, respectively.

law dependence of $\lambda_1(\varepsilon) \propto |\varepsilon|^{3/2}$. This dependence however disappears if we measure the instability exponent in the unit of the *dynamical time* t_D , which is customarily defined for SGS as proportional to the inverse square root of the density $t_D \propto (G\rho)^{-1/2}$. So, for SGS, we find that the adimensional chaoticity parameter γ_1 is constant and does not depends on the amount of binding energy in the system.

From the comparison of the two figures the peculiarity of SGS emerges clearly: *they behave, at any binding (specific) energy, as standard many body systems above the strong stochasticity threshold.*

This not only confirms that bound collisionless SGS are characterized by a dynamical instability proceeding at a very fast rate¹⁵, but also suggests that they possess the strongest statistical properties, in analogy with those of *standard* dynamical systems in the regime of fully developed stochasticity.

Once realized this, we are allowed to invoke the *Chaotic hypothesis* outlined in the Introduction, and to build a Statistical Mechanics for SGS.

3. Chaos and Statistical properties of Dynamics.

If the dynamics of a MDoF system possess strong enough stochastic properties, the Chaotic hypothesis (34) asserts that the averages of phase space functions can be computed as if the system were a transitive Anosov flow. From a physical viewpoint, this amounts to say that the system attains the most probable macroscopic equilibrium state compatible with the global constraints imposed at the initial time¹⁶.

¹⁵ This is indeed a very well known fact: it was discovered already by Miller in the '60s, when he performed the first pioneering numerical simulations of N-body gravitational systems and has been confirmed many times also in the last years.

¹⁶ These constraints are essentially the values of the possible conserved quantities, like energy, linear and angular momentum and so on.

So, if we are able to find this equilibrium state, we can compute all the averages using the invariant measure over the stationary distribution. The analogy with Anosov systems says even more, it gives also an indication about the *rates* governing the approach to the equilibrium state, which should be exponential if the manifold is *compact*.

For most MDoF systems these equilibrium (or stationary non equilibrium) states coincide either with the canonical or microcanonical ensemble, according to the system being in contact with a thermal bath or isolated, respectively. If the interaction potential governing the internal dynamics satisfies the conditions outlined in the Introduction, then the theorems of Statistical Mechanics entail that the descriptions based on the two ensembles agree, apart from terms which vanish as $1/N$ in the *thermodynamic limit*, *i.e.*, when the number of particles, the spatial volume and the energy of the system grow indefinitely, keeping however constant their ratios:

$$N \rightarrow \infty ; \quad V \rightarrow \infty ; \quad \mathcal{E} \rightarrow \infty \quad (16)$$

with

$$n \doteq \frac{N}{V} = \text{constant} ; \quad \varepsilon \doteq \frac{\mathcal{E}}{N} = \text{constant} . \quad (17)$$

3.1. *Obstacles TO A STATISTICAL MECHANICS FOR GRAVITATING SYSTEMS.*

From the above discussion, one amongst the many difficulties for a Statistical Mechanics of *gravitationally bound* SGS emerges immediately: it is not possible to perform the thermodynamic limit complying with the above requirements.

From the expression of the gravitational N-body Hamiltonian, equation (1), it is immediate to realize that keeping constant the density n implies that the specific energy grows without limit, $\varepsilon \propto N^{2/3}$. This problem derives obviously from the fact that, for *bound* SGS, the number of bodies (or better, the total mass), the energy, and the *spatial volume* cannot be fixed independently from each other, as it is instead possible for *standard* laboratory systems.

It is usually stated that that the one just mentioned is only the most evident difficulty and that another one, in principle not less relevant, is related to the fact that *the most probable state* for the gravitational N-body system cannot be determined, because *formally*, the measure of the phase space volume accessible for a given total energy is virtually infinite.

However, if we are not interested in the properties of the abstract, eternal, *final* equilibrium state, but rather to determine the quasi stationary states whose secular evolution proceeds on time scales much

longer than any dynamical process¹⁷, the second of the above mentioned difficulties can be surmounted, as we will show, in a (hopefully) self consistent way.

The remarkable fact is that, to overcome this obstacle, we do not need to avoid or mask the peculiarities originating it, but rather we succeed to incorporate their implications, to outline a hierarchical evolutive path, able to describe the succession of quasi-equilibria of stellar systems, covering a range of timescales spanning from those of the order of the virialization (or Violent Relaxation) up to those related to the formation of a collapsed-core-expanding-halo structure. As long as this hierarchy of timescales is covered, dynamical processes involving finer and finer spatial scales take place and this evolution is actually endless, as the *final* equilibrium state can be attained only on infinite times, and just for this, does not bear any direct relevance to *real* astronomical objects, once verified that the secular evolution of the metastable states proceeds on timescales longer than the stellar systems lifetime.

From the above discussion, it follows that the sole serious difficulty for a *standard* statistical description of N-body SGS is related to the non existence of the *standard* Thermodynamic limit. As we will discuss below, this is not a real hindrance for a self consistent Thermodynamics of SGS, as it is well known that many other *laboratory* systems¹⁸, characterized by non-extensive interactions, do not possess a well defined Thermodynamic limit¹⁹ and, nevertheless their macroscopic or collective behaviour can be described coherently using a suitable thermodynamic formalism which take into account their peculiarities.

The analysis below shows that a conceptually coherent framework can be settled to describe also SGS, certainly intrinsically even more peculiar than the *small systems* outlined above, but nevertheless suitable for a self consistent Thermodynamics.

3.2. DYNAMICAL SINGULARITIES AND STATISTICAL MECHANICS.

The above scenario owns also a dynamical counterpart which can be easily understood within the Geometro-dynamical framework. With-

¹⁷ And, from the astrophysical viewpoint, also much longer than the lifetimes of stellar systems to which we plan to apply at the end all these theoretical considerations.

¹⁸ The so called "*small systems*", see below.

¹⁹ Rather, the point is that, loosely speaking, the behaviour predicted in the T.L. does not bear any resemblance with the phenomenology observed in real, finite size, systems. And this should not to be considered surprising, because, for those systems, some *peculiarities* come just from the finiteness and disappear when the size exceeds the range of the interaction.

out going too much into details, we remark that the convergence of time averages to phase space values of geometric quantities (e.g., the curvature) proceeds in the case of SGS on timescales which strongly depend on the spatial resolution over which we compare the averages themselves. That is, the time required for the time averages (and fluctuations around them) to converge to the *static* averages is very sensitive to the spatial resolution used to compute the latter. Incidentally, we remark that the ergodicity time for the *Ricci curvature* is infinite if the mathematical definition of the Newtonian interaction is assumed. It becomes viceversa a well defined, finite, quantity, if the physical form of the interaction is considered. It must be stressed that even in this respect, as the spatial resolution increases, the ergodicity timescales of geometrodynamical quantities grow up, exactly at the same rate of the times associated to the relaxation of statistical quantities.

3.2.1. *Mathematical vs. Physical peculiarities of gravitational interaction.*

We conclude this section listing the more relevant peculiarities of the Newtonian interaction outlining also which, if any, amongst them could constitute a real obstacle to a functional statistical mechanical foundation of a Thermodynamics for SGS.

Singular interaction at short distances: The most evident singularity of the gravitational interaction is obviously related to the divergency of the potential when two particles get closer and closer. Although clearly of utmost importance for the single particle dynamics and for the evolution of *few body* systems, the global and statistical relevance of close encounters is practically negligible over any finite timescale, as the frequency of high energetic close encounters is vanishingly small and the effect on the global quantities irrelevant, compared with that of the overall dynamics. Then, even if it is physically plausible²⁰, from a statistical viewpoint there is no need of any *softening* of the potential.

Non confining nature of the potential and divergence of the phase-space volume: The configuration space of a N-body bound SGS obviously diverges, and in principle very badly, indeed as

$$\Gamma_{\mathbf{q}} \sim \lim_{V \rightarrow \infty} V^{N-2}.$$

This is due to the fact that Newtonian interaction is unable to confine, asymptotically, inside a finite volume more than two

²⁰ And numerically very useful, especially if low order efficient symplectic integrators are used, in order to guarantee the faithful preservation of the Hamiltonian properties of the flow.

bodies. If the mathematical N-body system is considered, an analogous divergence comes out also from the volume in the momentum space. However, as long as $\log N \gg 1$, the characteristic time for the evaporation is considerably longer than any dynamical timescale, and the evolution is only secularly influenced by the slow evaporation of stars, which, incidentally, causes the remaining system being more and more bound, so inhibiting further escapes.

Unscreened, long-range nature of the interaction: It is the singularity which most affects the statistical and thermodynamics properties of SGS. This is in part obvious as it concerns the dynamics over all the spatial scales, and consequently, according to the general discussion above, every timescale. The long range nature of the interaction reflects on the efficiency of collective phenomena, among which the Violent Relaxation is surely the most popular but it is not the only one. The absence of any effective screening mechanism entails that gravitationally bound systems do not own any characteristic length, and reflects on the scale invariance of most features. From statistical and thermodynamical viewpoints the most far reaching consequences reflect in the following points:

- N-body systems are not *stable* in the sense of *Rigorous Statistical Mechanics* (1): given any number of particles N , the binding energy per particle can increase without any upper bound (*i.e.* the energy does not have a *lower* bound). In particular, the binding energy of the *fundamental state* of a system scales at least as N^2 . That the lack of stability property is related to the long range nature of the potential and it is unaffected from the singularity at short distances emerges clearly from the discussion above and from many numerical simulations performed recently (44). Indeed, the introduction of a softening of the potential is able to introduce an upper limit for the binding energy of the system, but is unable to *cure* the lack of the *stability* property²¹.
- From the previous point, it follows that the internal energy of a system scales as N^2 , and this remains true for most thermodynamic potentials. As the extensivity of the latter is a necessary condition for the proof of the ensemble equivalence, the long range nature of the Newtonian interaction is the main cause of the peculiarity of Gravitational Thermodynamics.

²¹ Even the most *radical* modification of the short range interaction, *i.e.*, the introduction of an hard-core repulsive potential at short distances, does not prevent the divergence of the binding energy per particle in the thermodynamic limit, though being able to make it finite at any finite N .

From the above discussion it emerges clearly that is the non-extensivity of potential energy (and, by the virial theorem, also of the kinetic and total energies) to imply the non equivalence of statistical ensembles, and this requires, in turn, a deeper analysis in order to understand which ensemble, if any, is best suited to describe SGS.

4. From Statistical Mechanics to Microcanonical Thermodynamics of self gravitating systems.

Up to now we worried about the *rigorous theorems* concerning the equivalence of ensembles, the scaling of macroscopic quantities in the thermodynamic limit and so on. However, as emphasized in previous sections, in recent years it has realized that the issue of the inequivalence of statistical ensembles emerges whenever the scale of the dominating interaction is comparable with the spatial extension of the system under study. It is easy to understand that in such circumstances the interaction energy ceases to be an extensive quantity and, if some *equipartition property* holds, the same is true also for kinetic and total energies. However, many of these so-called *small systems* (15) possess a well defined, though somewhat peculiar, thermodynamic behaviour. In these systems it is not difficult to encounter regions in the space of thermodynamic parameters where the specific heats become negative. When described within the microcanonical ensemble, all these peculiar phenomena do not originate any *paradox*, which appears instead as soon as a *canonical ensemble* approach is attempted. The best that can be done, sometimes, within this framework, is to describe the phenomenology inside these regions as a phase transition, though *true* phase transitions themselves are *rigorously* defined only in the Thermodynamic limit, which we know to be a rather meaningless limit just when the system properties are not extensive.

4.1. ENSEMBLE INEQUIVALENCE.

As discussed in the previous sections, the non extensive nature of the energy entails, among others, the inequivalence of statistical ensembles. In order to find out what among the classical ensembles is best suited to describe SGS we first observe that, if we consider *isolated* N-body systems, nothing suggests to suppose them in contact with any heat bath. Going a little more below the surface, we observe more concretely that if we require, according to Boltzmann, that the ensemble fulfils the *orthodic* property (see (2), §5), then neither the canonical nor the grand-canonical ensembles can be used to describe gravitationally

bound N-body systems. Heuristically, this simply because both these ensembles fix the *Temperature* as a constant parameter of the distribution, assuming implicitly that this quantity is constant throughout every system of the ensemble.

To this somewhat formal point, we have to add the already quoted, and physically most relevant, incongruence met by the canonical (and also grand-canonical) approach: SGS are characterized by negative heat capacities, whereas equation (3) shows that within the canonical ensemble the heat capacity is positive by definition. Moreover, the non extensivity of potential, kinetic and total energies entails the analogous non extensivity of canonical and grand-canonical thermodynamics potentials, as the Helmholtz free energy F or the Gibbs potential G . These features not only imply the non equivalence of ensembles but also indicate that even the definition of averages (and even more of fluctuations) of thermodynamic variables becomes problematic.

Up to this point we analyzed the problems affecting the canonical and grandcanonical descriptions. Now we turn to ask how the peculiarities of gravitational interaction reflect also on the microcanonical description.

Before to proceed, we think that this seems to be the appropriate point to recall a general statement: the *strong* chaotic properties of dynamics, the Poincaré-Fermi theorem, the general theorems of Ergodic Theory, all focus on the relationships between the stochastic properties of motions and the occurrence of a uniform diffusion of initially concentrated phase space volumes over the whole region accessible to motion, compatibly with the integrals of motion allowed by the system dynamics. If such a spreading process in phase space is efficient enough, then, those general theorems, also according to the *Chaotic Hypothesis*, assert that the time averages of dynamical variables coincide with the phase averages computed over the region of phase space where motions take place transitively. If we are dealing with Hamiltonian systems, this region is the constant energy surface²² Σ_E . This amounts to say that the *strong enough* stochastic properties of Dynamics of MDoF Hamiltonians, naturally lead to a *microcanonical* statistical description.

The task of the powerful formal apparatus of *rigorous* Statistical Mechanics is then to *prove* the equivalence of ensembles. In case of success, then we can use the more *tractable* expression of canonical

²² Possibly taking also into account the conservation of other global integrals, like the position and linear momentum of the center of mass of the system and the total angular momentum if only central forces are present. This reduces the dimensionality of Σ_E from $6N - 1$ to $6N - 10$, which, for $\log N \gg 1$, does not modify obviously the macroscopic behaviour.

or grand-canonical ensembles to describe the macroscopic properties and the Thermodynamics of our system. If the equivalence does not hold, then the sole Statistical description allowed is the one compatible with the global constraints on our system: if it is isolated, then only a microcanonical description is *legitimate*.

From the above discussion, it is clear that the use of *microcanonical description* is justified, provided that no internal inconsistencies occur; and that the inequivalence of ensembles, if ever, should suggest a critical reconsideration of the results obtained using a canonical or grand-canonical description.

It is easy to see that the only trouble comes from the formal unboundness of the phase space volume accessible at a given constant energy. Nevertheless, if we are interested in the finite time quasi equilibrium states, we can take the argumentations discussed in the previous sections to emphasize that, at any finite time the *effective* phase space volume accessible is well defined in terms of the global quantities describing the state of the system (energy per particle, specific angular momentum, etc.) and it is finite²³.

So, for any finite scale of time, we are able to derive an *effective* expression for the microcanonical entropy, secularly evolving, and able to take into account the increasing effects of the phenomena occurring on smaller and smaller spatial scales, and, correspondingly, on longer timescales.

Starting from the Hamiltonian of the system (1), given a value for the total energy of a *bound* SGS²⁴, $\mathcal{E} < 0$, we can define a set of *macroscopically averaged* quantities:

the energy per particle:

$$\varepsilon \doteq \frac{\mathcal{E}}{N} \leq 0 , \quad (18)$$

²³ It is easy to verify that even in the presence of escaping particles or taking into account the occurrence of close encounters, the contribution of these events to the actual phase-space volume is negligible as long as $\log N \gg 1$ for times of $t \geq \mathcal{O}(Nt_D)$.

²⁴ It is possible to perform a statistical mechanical treatment even for loosely bound systems or systems in interaction with some background gravitational field. This leads to a very interesting phenomenology, indicating the occurrence of critical phenomena in 3D-gravitational systems. The details of this work are described in (44).

the virial ratio:

$$\mathcal{Q}(t) \doteq \frac{-2K(t)}{U(t)} \equiv \frac{\sum_{i=1}^N \frac{p_i^2}{m_i}}{\frac{1}{2} \sum_{i=1}^N \sum_{j \neq i} \frac{m_i m_j}{r_{ij}}} , \quad (19)$$

where the indication of the time dependence of the kinetic and potential energies, $K(t)$ and $U(t)$, respectively, is a shortcut to recall that their values depend on the state of the system at time t ;

the inertial radius:

$$R_i(t) \doteq \left[\frac{\sum_{i=1}^N m_i r_i^2}{\sum_{i=1}^N m_i} \right]^{1/2} , \quad (20)$$

the harmonic (or virial) radius:

$$R_g(t) \doteq \frac{\left(\sum_{i=1}^N m_i \right)^2}{\sum_{i=1}^N \sum_{j \neq i} \frac{m_i m_j}{r_{ij}}} , \quad (21)$$

and the ratio among them:

$$\alpha(t) \doteq \frac{R_i(t)}{R_g(t)} \quad (22)$$

In the simplifying hypothesis that the mass spectrum is nearly flat, so that the average mass $\bar{m} \doteq \langle m_i \rangle_i$ represents the typical mass of the system²⁵, it is a simple computation to obtain an expression for the microcanonical entropy of the system, which reads

$$S \equiv k_B \sigma(N, \varepsilon, \mathcal{Q}, \alpha, \bar{m}) , \quad (23)$$

where

$$\sigma = \frac{3N}{2} \ln \left[\left(\frac{G^2 N^2 \bar{m}^5}{-2\varepsilon} \right) \alpha^2(t) \mathcal{Q}(t) [2 - \mathcal{Q}(t)] \right] \quad (24)$$

²⁵ When a very steep spectrum of masses is allowed, the computations are only a little more involved, but the results presented below are unchanged. In addition it is possible to obtain also the dependence of the entropy on the mass spectrum.

and it is obviously understood that the also the varying quantities $\alpha(t)$ and $Q(t)$ depends on time only through the actual state of the system.

The above argument can be extended to take into account other details about the state of the system, but the simplest derivation sketched here is sufficient to emphasize a number of relevant points.

It is evident that the microcanonical entropy (24) gives correctly the equilibrium conditions related to variations of the parameters on which it depends.

However, the most striking result is that it turns out to be *an extensive quantity*, being proportional to the number of particles²⁶.

Another very important feature of this entropy is that it is possible to derive from it easily all the relevant thermodynamic quantities, which agree with the phenomenological expressions obtained at a rather heuristic level.

The most important is the *microcanonical temperature* which, coherently with the statistical description, is clearly only an average quantity for the systems of the microcanonical ensemble²⁷

$$\frac{1}{T} \doteq \left(\frac{\partial S}{\partial \mathcal{E}} \right)_N \equiv \frac{1}{N} \left(\frac{\partial S}{\partial \varepsilon} \right)_N \quad (25)$$

which, using (24), gives, coherently with the fact that bound systems have negative energy and positive temperature,

$$\varepsilon = -\frac{3}{2}k_B T . \quad (26)$$

Consequently, a negative heat capacity follows:

$$C_V \doteq \left(\frac{\partial \mathcal{E}}{\partial T} \right)_N = -\frac{3}{2}Nk_B ; \quad (27)$$

that is, a *negative specific heat*, $c_V = -3/2k_B$.

The surprising features of the entropy (24), defined in such a straightforward way, do not limit to single out the thermodynamic properties of the system, but allows also to determine the dynamical evolution of the global quantities. It is immediate to see, for example, that, as a function of the virial ratio, the entropy (24) *is maximized when $Q = 1$* :

$$\frac{\partial \sigma}{\partial Q} = \frac{2(1 - Q)}{Q(2 - Q)} \quad (28)$$

²⁶ A very general approach to the definition of suitable microcanonical entropies for generic MDoF systems is discussed in (45).

²⁷ According to what stated above, in the derivation of the microcanonical thermodynamic relations, we take into account that, for bound SGS, the energy, the number of bodies and the volume are not independent quantities.

Figure 4. Probability distribution for the values of the virial ratio, determined according to the definition of the microcanonical entropy, equation (24), for bound self gravitating systems with moderately large number of bodies: from the broader to the narrower distributions it is $N = 10, 50, 200, 1000$. It is well evident that, as the number of bodies increases, the overwhelming fraction of $\Sigma_{\mathcal{E}}$ is occupied by systems in virial equilibrium.

That is, during the approach towards the virial equilibrium the average entropy of the system increases to the maximum value allowed by the values of the other parameters. Equivalently, it can be said that, starting from an initial state out of the virial equilibrium, entails a rapid relaxation to the maximum entropy state compatible with the given energy. That most of the volume of phase space of N-body systems is occupied by configurations in virial equilibrium is evident from figure 4, where the probability distribution of the $\mathcal{Q}$ values is shown to be strongly peaked around the equilibrium value already for moderately large N-values.

We think that it is very impressive to see how such a definition of *finite time entropy* is able to predict even the dynamical states maximizing locally (in time) the entropy.

As a further evidence of this property, we observe that it is also possible to predict (and also to numerically verify), that the secular dynamical evolution of a gravitationally bound system leads the ratio $\alpha(t)$ to an average steady growth, over which small quasi periodic modulations are imposed. The expression derived above for the microcanonical entropy correctly describes also this secular dynamical evolution through locally stationary states of slowly increasing entropy!

$$\frac{\partial \sigma}{\partial \alpha} \propto \frac{N}{\alpha} > 0 . \quad (29)$$

This means that the entropy will increase along with the growth of the parameter $\alpha(t)$ (because, supposing fixed all the other parameters, we have $\dot{\sigma} \propto N\dot{\alpha}/\alpha$). Well, it is easy to see, even analytically, that the development of a core-halo structure implies an increase of $\alpha(t)$. The relationship between σ and α however indicates that the entropy growth rate, though remaining positive, slows down as α steadily increases.

Adding a further comment to the results above, we remark that, while the apparent singularity of the entropy when $\varepsilon \rightarrow 0^-$ is fictitious, as, in this limit it is also $\mathcal{Q} \rightarrow 2^-$, the singularity for $\mathcal{Q} \rightarrow 0^+$ is instead real, and again very enlightening on the sensible meaning of the microcanonical entropy here presented. Indeed the above limit describes a *completely frozen* state of the system, that is, a given configuration with all the bodies having zero velocity. The probability of such a state

is zero for any $\varepsilon < 0$, and correctly its entropy diverges. Incidentally we notice that, if $\varepsilon \rightarrow 0^-$, then a *frozen* state become possible, and accordingly σ can be made finite.

We emphasize again how the present definition of microcanonical entropy is effective in taking into account the secular evolution of the system, through the full hierarchy of timescales associated to the singular dynamical processes occurring in the asymptotic evolution of SGS. It is also remarkable that, if we consider a somewhat *regularized* gravitational N-body system, where the divergence at short distance is removed (e.g., through a usual softening procedure, for example) and the unboundness of configuration space accessible is avoided, introducing a kind of *cosmic vessel*, then the microcanonical entropy will still entail the secular evolution, but will be in this case able to single out an *end state* that is now well defined and attainable in a finite time! We stress that this is not at all a trivial result, because the *regularization* procedure described above does not modify the statistical and thermodynamic peculiarities of the system, which remains nonextensive and for which the inequivalence of ensembles remains true, as long as the system is gravitationally bound. In particular, such a regularized system will keep a negative heat capacity in a wide range of specific energies and the canonical ensemble description permains unable to grasp even the qualitative features of its Thermodynamics (44)!

5. Comments and Conclusions.

We presented an *effective microcanonical entropy* able to describe correctly not only the thermodynamic properties of SGS, but also to outline the sequence of secular evolutionary processes leading the systems towards states of higher and higher entropy.

The dynamical and statistical justifications of a microcanonical description of gravitationally bound N-body systems have been derived and discussed on the grounds of analytical and numerical evidences of their *strongly chaotic properties*, which allow the assumption of validity of the *Chaotic Hypothesis* of Gallavotti and Cohen. The results obtained within the *Geometro-dynamical Approach to Chaos* gave further support to the reliability of this assumption.

The peculiar dynamical properties of gravitational interaction, the breakdown of the extensivity and the consequent inequivalence of classical statistical ensembles, moreover, rule out any reliable canonical or grand-canonical description of the Statistical Mechanics and Thermodynamics of *gravitationally bound* N-body systems.

It can be shown that the microcanonical entropy is the sole thermodynamic potential which keeps the extensivity at any finite time even for the non stable Newtonian interaction potential, while all thermodynamic potentials defined within the canonical or grand-canonical ensembles lose the extensivity property and consequently lead to unclear, unuseful or wrong thermodynamic relations. A simple sketch of the above assertions follows directly from the definitions of the canonical and grand-canonical potentials:

Helmholtz free energy.

In the canonical ensemble, given the internal energy

$$U \equiv \langle H \rangle_\beta \cong \mathcal{E} , \quad (30)$$

where average is performed on the canonical ensemble and $\mathcal{E}$ is instead the fixed energy of the *corresponding microcanonical ensemble*²⁸; the Helmholtz free energy is related to it by the relation

$$F = U - TS . \quad (31)$$

From the results obtained above, we see clearly that, for a non extensive system (be it a SGS or a *small system*) it is $F \propto N^2$; that is, F is not more an extensive quantity, although S it is²⁹.

Gibbs potential.

In the grand-canonical ensemble, the appropriate potential is given by the Gibbs function, whose thermodynamic relationship with F is given by

$$G = F + PV$$

and, as it is easy to verify, both the terms of the sum are, for a *bound* SGS, proportional to N^2 .

If a *physically reasonable regularization* of gravitational interaction, which does not modify its characteristic peculiarities, like the non extensivity and the *non stability*, is introduced, the proposed microcanonical entropy keeps completely its validity and moreover becomes able to single out not only the hierarchy of secularly evolving meta-equilibrium states, in analogy with the non regularized case, but also

²⁸ We are using the relations which are valid when there is the ensemble equivalence. Here we use them to further demonstrate the difficulties of canonical description!

²⁹ At this point we notice that the above derived expression for the temperature, equation (26), gives, $T \propto \varepsilon$. For non extensive systems this entails that also the product TS is proportional to N^2 , coherently with all the above discussions.

to determine a non singular *final* equilibrium state, which was only asymptotically reachable prior the regularization.

In addition, the above regularization allows for a detailed study of the critical phenomena occurring in SGS (44), able to show the occurrence of phase transitions in gravitational N-body systems and also to confirm the doubtful reliability of a canonical description of the Statistical Mechanics and Thermodynamics of Gravity .

The above discussion points out that the peculiarity of the collective properties of gravitational systems depend largely on the long range attractive nature of the interaction and it is almost completely independent on the formal singular behaviour at short distances. Taking into account that the latter is only a formal singularity, we can safely ignore it also on the grounds of the results showing its complete irrelevance for the statistical properties of SGS. The details on the way of coping with the dynamical consequences of the presence of the singularity affect the evolution of macroscopic quantities only on timescale which are longer than the binary relaxation time, in some cases even exponentially larger!

5.1. FURTHER COMMENTS AND FUTURE DEVELOPMENTS.

A further point deserves to be remarked: it is undeniable that the *limits* on which most of the *standard* Statistical Mechanical and Thermodynamic frameworks rely are of outmost relevance for the comprehension of the behaviour of most macroscopic systems. For the overwhelming largest class of MDoF systems, the thermodynamic limit allows to understand the basic and usual properties of macroscopic matter and even the rich and complex phenomenology of critical phenomena. At the same time, theorems assuring the existence of an invariant distribution are exceptionally important to single out the properties of the final equilibrium states of most macroscopic systems.

Nevertheless, it is also evident from the considerations here presented, that the operation of taking these limits (*i.e.*, the thermodynamic limit and the infinite time limit), assigning to them the role of selection criteria in order to establish whether a *Thermodynamics* for a given class of systems can be constructed or not, can hinder in some cases the possibility of obtaining alternative statistical and thermodynamic description of MDoF systems which are certainly *singular* in some respect, but possess notwithstanding well defined, though peculiar, thermodynamic properties, in general for many of them, or at least over any physically reasonable finite interval of time, as it happens

for SGS.

The outcomes presented above are enlightening also for the critical analysis of the results based in some way on the assumption of a canonical equilibrium of gravitational N-body systems. In particular, we claim that many results based on the maximization of entropy like quantities constructed using the single particle distribution function have to be reconsidered.

In conclusion, despite the *rigorous theorems*, we have shown that a thermodynamic description of gravitational N-body systems is not only feasible, but also suggests a new approach to *peculiar systems* for which the standard paradigms of *rigorous* Statistical Mechanics do not apply.

We plan to further develop the line of research here proposed in order to make it possible to obtain indications on the detailed evolution of SGS on the basis of thermodynamic arguments alone.

In order to attain this goal it is necessary to have more informations on the detailed energy distribution of N-body systems and for this the use of extensive numerical simulations is required.

Acknowledgements

The work of P.C. is partially supported by C.S.S. under the initiative n.2000A:MPDS.

References

- (1) D.Ruelle, *Statistical Mechanics: rigorous results* - Addison-Wesley Publishing Company, 1969.
- (2) G.Gallavotti, *Meccanica Statistica, Trattatello* - Quaderni del C.N.R.-G.N.F.M., n.50, 1995, *in italian*.
- (3) S.Chandrasekhar, *Principles of Stellar Dynamics* - Dover Publ. (reprint), 1969.
- (4) T.Padmanabhan, *Phys. Rep.* **188**, 285, 1990.
- (5) W.C.Saslaw, *Gravitational Physics of Stellar and Galactic Systems* - Cambridge Univ. Press, 1985.
- (6) J.Binney, S.Tremaine, *Galactic Dynamics* - Princeton Univ. Press, 1987.
- (7) D.Lynden-Bell, *M.N.R.A.S.* **136**, 101, 1967.
- (8) W.C.Saslaw, *M.N.R.A.S.* **150**, 299, 1970.
- (9) F.H.Shu, *Ap.J.* **225**, 83, 1978; - *Ap.J.* **316**, 502, 1987.
J.Madsen, *Ap.J.* **316**, 497, 1987.
J.Hiort, J.Madsen, *M.N.R.A.S.* **265**, 237, 1993.
- (10) P.D.Louis, O.E.Gerhard, *M.N.R.A.S.* **233**, 337, 1988.
P.Mineau, M.R.Feix, J.L.Rouet, *Astron.Astroph.* **228**, 344, 1990.

- (11) W.Thirring, Z.Physik **235**, 339, 1970.
P.Hertel, W.Thirring, Ann. of Phys. **63**, 520, 1971.
- (12) V.A.Antonov, Vest.Leningr.Univ. **7**, 135, 1962.
- (13) D.Lynden-Bell, R.Wood M.N.R.A.S. **136**, 101, 1967.
- (14) K.Huang, *Statistical Mechanics* - John Wiley & sons, NY, 1964.
- (15) D.H.E.Gross, Phys.Rep. **279**, 119, 1997.
- (16) D.Lynden-Bell, R.M.Lynden-Bell, M.N.R.A.S. **181** 405 1977.
- (17) Ph.Chomaz, V.Duflot, F.Gulminelli, Phys.Rev.Lett **85**, 3587, 2000.
- (18) T.Dauxois, P.Holdsworth, S.Ruffo, Eur.Phys.J. **B 16**, 659, 2000.
- (19) M.Pettini, Phys.Rev. **E 47**, 828, 1993;
M.Cerruti-Sola, M.Pettini, Phys.Rev. **E 51**, 53, 1995;
M.Cerruti-Sola, M.Pettini, Phys.Rev. **E 53**, 179, 1996.
- (20) P.Cipriani, *Ph.D.Thesis*, Univ. of Rome “La Sapienza”, 1994 (in italian);
P.Cipriani, G.Pucacco, N.Cimento **B 109**, 325, 1994;
P.Cipriani, G.Pucacco, in *Ergodic Concepts in Stellar Dynamics*, (V.G. Gurzadyan - D. Pfenniger eds.), Lecture Notes in Physics:vol.430, Springer-Verlag, 163, 1994.
- (21) L.Casetti, M.Pettini, Phys.Rev. **E 48**, 4320, 1993;
L.Casetti, R.Livi, M.Pettini, Phys.Rev.Lett. **74**, 375, 1995.
- (22) P.Cipriani, M.Di Bari, Plan.Sp.Sci. **46**, 1499, 1998;
M.Di Bari, P.Cipriani, Plan.Sp.Sci. **46**, 1534, 1998.
- (23) P.Cipriani, M.Di Bari, Phys.Rev.Lett. **81**, 5532, 1998;
M.Di Bari et al. Phys.Rev. **E 55**, 6448, 1997.
- (24) L.Casetti, M.Pettini, E.G.D.Cohen, Phys.Rep. **337**, 237, 2000.
- (25) D.Pfenniger, Astron.Astroph. **165**, 74, 1986;
H.E.Kandrup, M.N.R.A.S. **299**, 1139, 1998;
D.Merritt, P.A.S.P. **111**, 129, 1999.
- (26) S.Tremaine, M.Henon, D.Lynden-Bell, M.N.R.A.S. **219**, 285, 1986;
S.D.Mathur, M.N.R.A.S. **231**, 367, 1988;
H.E.Kandrup, M.N.R.A.S. **225**, 995, 1987.
- (27) H.Poincaré, *Les Méthodes nouvelles de la Mécanique Céleste*, Paris, 1892.
- (28) M.Hénon, C.Heiles, Astron.J. **69**, 73, 1964.
- (29) V.I.Arnold, *Mathematical Methods of Classical Mechanics* - Springer-Verlag, 1980;
A.Katok, B.Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems* - Cambridge Univ. Press, 1995;
Ya.G.Sinai, (Ed.) *Dynamical Systems* - World Scientific, 1991.
- (30) G.Benettin, in *Non linear evolution and Chaotic phenomena*, (G.Gallavotti - P.F. Zweifel eds.), Plenum, p.121, 1988.
- (31) L.Galgani, in *Non linear evolution and Chaotic phenomena*, (G.Gallavotti - P.F. Zweifel eds.), Plenum, p.147, 1988.
- (32) E.Fermi, in *Collected Papers of E. Fermi vol.II*, (E.Segré ed.), Univ. of Chicago Press, 1965.
- (33) M.Falcioni, S.Isola, A.Vulpiani, Phys.Lett. **A 144**, 341, 1990;
M.Pettini, M.Landolfi, Phys.Rev. **A 41**, 768, 1990;
T.Tèl, P.Gaspard, G.Nicolis (eds.) in *Chaos Focus issue on chaos and irreversibility* **8**, 309, 1998.
- (34) G.Gallavotti, E.G.D.Cohen, Phys.Rev.Lett. **74**, 2694, 1995;
G.Gallavotti, E.G.D.Cohen, J.Stat.Phys. **80**, 931, 1995.
- (35) J.L.Lebowitz, O.Penrose, Physics Today, 23 (February 1973).
- (36) D.V.Anosov, Proc.Steklov Inst. Math. **90**, 1, 1967.

- (37) N.S.Krylov, *Works on Foundations on Statistical Physics*- Princeton Univ. Press, 1979.
- (38) J.L.Synge, Phil. Trans. **A 226**, 31, 1926.
- (39) J.Berndt, L.Vanhecke, Differ.Geom.& Appl. **2**, 57, 1992.
- (40) L.D.Landau, E.M.Lifsits, *Mechanics* (3^{rd} edition), Pergamon, 1976.
- (41) P.Cipriani, Il N.Cimento **B 115**, 1109, 2000.
- (42) N.G.Van Kampen, Phys.Rep., **24**, 171, 1976.
- (43) G.Benettin, L.Galgani, J.M.Strelcyn, Phys.Rev. **A 14**, 2338, 1976.
- (44) M.Cerruti-Sola, P.Cipriani, M.Pettini, M.N.R.A.S., *submitted*, 2000.
- (45) P.Cipriani, to be submitted to Phys.Rev. E, (2001).







